

Galaxy Evolution tool: Construction and Applications

Yeshe Fenner and Brad K. Gibson

Centre for Astrophysics & Supercomputing, Swinburne University, Australia

Marco Limongi

Osservatorio Astronomico di Roma

Keywords: Galaxy: evolution, nucleosynthesis, abundances

Galaxy Evolution tool (GEtool) is a software package currently being developed to self-consistently model the chemical and spectral evolution of disk galaxies. GEtool will soon be available to the community through a web-based interface¹ that will enable users to predict observable properties of model galaxies such as colours, spectral gradients, Lick indices and elemental abundances. We present an application of the code to a dual accretion phase model of the Milky Way in order to assess the role of massive stars in the chemical composition of the Galaxy. A updated set of stellar yields, covering a range of stellar initial masses and metallicities ($13 < m/M_{\odot} < 80$ and $Z/Z_{\odot} = 0, 10^{-3}, 1$), has recently been calculated by Limongi et al (2001, in prep) (LSC01). This poster presents results from the first chemical evolution model to incorporate these new stellar yields.

The age-metallicity relation, G-dwarf distribution and evolution of abundance ratios predicted using the sets of yields from LSC01 and Woosley & Weaver (1995, ApJS, 101, 181) (WW95) are compared with observations. Stellar yields are one of the most important ingredients in Galactic chemical evolution models, yet the ejected mass of iron-peak elements is very uncertain due to the edge of the iron core being so close to the mass cut. Further uncertainties are introduced by the possibility that much of the synthesised iron in higher mass stars falls back to core. Since there are few supernova observations with which to directly infer the iron ejecta, we compared the predicted and observed trend of key abundance patterns to indirectly constrain iron yields. We found that the trend of $[O/Fe]$ and $[Mg/Fe]$ vs $[Fe/H]$ inferred from recent measurements cannot be recovered if the iron yield of massive stars increases with initial mass, as in WW95 model. Instead, the observational constraints are better satisfied if we assume the iron yield decreases for initial masses greater than $\sim 25 M_{\odot}$ (see Figure 1 and Figure 2.a.). Both models shown in Figure 2.b. reproduce the observed G-dwarf distribution in the solar neighbourhood.

¹ <http://astronomy.swin.edu.au/GEtool/>




Figure 1. The ejected mass of O (*left*) and Fe (*right*) as a function of main sequence mass from LSC01 (*grey lines*) and WW95 (*black lines*). The estimated mass of iron ejected from five observed SNe are shown. Two models have been proposed for SN1998bw.




Figure 2. *a.* O/Fe vs metallicity for stars are plotted for comparison with predictions from dual-infall chemical evolution models. Grey lines represent models using LSC01 yields while black lines depict results from WW95. Observational data are from Edvardsson et al. (1993, A&AS, 102, 603: *filled circles*), Israelian et al. (2001, ApJ, 551, 833: *crosses*), Boesgaard et al. (1999, ApJ, 117, 492: *triangles*) and Carretta et al. (2000, A&A, 356, 238: *diamonds*). *b.* The predicted G dwarf distribution from the two stellar yield models (*grey and black lines*) are shown against the observed metallicity distribution of G and K dwarfs in the solar neighbourhood (*histograms*).